

A Capture Problem in Brownian Motion and Eigenvalues of Spherical Domains

Jesse Ratzkin
University of Connecticut
ratzkin@math.uconn.edu

Andrejs Treibergs
University of Utah
treiberg@math.utah.edu

February 1, 2008

Abstract

We resolve a question of Bramson and Griffeath by showing that the expected capture time of four independent Brownian predators pursuing one Brownian prey on a line is finite. Our main tool is an eigenvalue estimate for a particular spherical domain, which we obtain by a coning construction and domain perturbation.

In this paper, we examine the expected capture time of a single Brownian prey pursued by n independent Brownian predators. All motion is restricted to a line. Bramson and Griffeath [BG] first considered this problem, and estimated the capture time in various circumstances. In particular, they showed that if at time $t = 0$ there are predators on both sides of the prey then the expected capture time is finite. For this reason, we will assume that the initial position of the prey is $x_0(0) = 1$ and the initial positions of the predators are $x_1(0) = \dots = x_n(0) = 0$. In this case, [BG] showed that the expected capture time is infinite for $n = 1, 2, 3$, and conjectured that it is finite for $n \geq 4$ (as indicated by simulations). Li and Shao [LS] showed that the expected capture time is finite for $n \geq 5$. Using a similar method, we resolve the remaining case by showing that the expected capture time is finite for $n = 4$.

Theorem 1 *Let $x_0(t)$ be pursued by $x_1(t), \dots, x_n(t)$, where $x_0, \dots, x_n$ are independent, standard Brownian motions on a line, $x_0(0) = 1$, and $x_1(0) = \dots = x_n(0) = 0$. Then the expected capture time is finite if and only if $n \geq 4$.*

One key difference between Li and Shao's method in [LS] and our method here is that they consider a difference process, while we do a geometric splitting in spherical polar coordinates to reduce the dimension of the problem. See Section 2 for more details on our dimension reduction.

One can reformulate the capture problem described above in terms of exit of a Brownian particle in $\mathbb{R}^{n+1}$ from a specific cone. We denote the position of the prey at time t as $x_0(t)$ and the position of the j th predator at time t as $x_j(t)$. By our choice of initial conditions, the initial position of the Brownian particle $x(t) = (x_0(t), x_1(t), \dots, x_n(t))$ is $x(0) = (1, 0, \dots, 0)$. The event of capture is then equivalent to the Brownian particle $x(t)$ first leaving the cone

$$\mathcal{C}_{n+1} := \{(x_0, x_1, \dots, x_n) \mid x_0 \geq x_j, j = 1, \dots, n\},$$

with $x(0) = (1, 0, \dots, 0)$, and so we must estimate the expected first exit time of a Brownian particle from the cone $\mathcal{C}_{n+1}$, with the starting position $(1, 0, \dots, 0)$.

DeBlassie [DB] developed the theory of estimating exit times for Brownian motion from cones in Euclidean space. Let $C = \{(r, \theta) \mid r \geq 0, \theta \in D \subset \mathbb{S}^n\}$ be the cone over a domain $D \subset \mathbb{S}^n$. Also let τ_x be the exit time from C of a Brownian particle with starting position x , and let $\mathbb{P}(\tau_x > t)$ be the probability that $\tau_x > t$. DeBlassie showed $\mathbb{P}(\tau_x > t) \sim c(x)t^{-a(n)}$, where

$$2a(n) = \left[\left(\frac{n-2}{2} \right)^2 + \lambda_1(D) \right]^{1/2} - \frac{n-1}{2}. \quad (1)$$

Here $\lambda_1(D)$ is the first Dirichlet eigenvalue of D . In the particular case we are interested in, the expected exit time of a Brownian particle from $\mathcal{C}_{n+1}$ is finite if and only if $a(n) > 1$, which reduces to

$$\lambda_1(\mathcal{D}_n) > 2n + 2, \quad (2)$$

where $\mathcal{D}_n = \mathcal{C}_{n+1} \cap \mathbb{S}^n$.

Our method for proving Theorem 1 is to estimate the first eigenvalue of $\mathcal{D}_n$ using the monotonicity property of eigenvalues, a coning construction, and domain perturbation. The rest of the paper proceeds as follows. In Section 1 we discuss the geometry of $\mathcal{C}_{n+1}$, $\mathcal{D}_n$, and related regions. Section 2 contains the separation of variables background one needs to estimate the expected capture time. For the reader's convenience, we also include the proof that the expected capture time is infinite for $n = 1, 2, 3$ predators in this section. We prove the $n = 4$ eigenvalue estimate in Section 3. Finally, we describe a numerical computation of $\lambda_1(\mathcal{D}_3)$ and a lower bound for $\lambda_1(\mathcal{D}_4)$ in Section 4.

ACKNOWLEDGEMENT: Davar Khoshnevisan told us about this problem. We thank him and Pedro Mendez for lending their ears and expertise in this project. A. T. thanks Frank Stenger for suggesting the algorithm of Section 4.

1 Geometry of the cone $\mathcal{C}$

The cone $\mathcal{C}_{n+1}$ and its spherical angle $\mathcal{D}_n$ have much symmetry. First observe that $\mathcal{C}_{n+1}$ contains the line $\mathcal{L}$ spanned by $(1, 1, \dots, 1)$; this is the line where all the inequalities $x_0 \geq x_j$, $j = 1, \dots, n$ are equalities. Thus we can split $\mathcal{C}_{n+1}$ as a sum of a line and lower dimensional cone

$$\mathcal{C}_{n+1} = \mathcal{L} \oplus \mathcal{V}_n,$$

where $\mathcal{V}_n = \mathcal{L}^\perp \cap \mathcal{C}_{n+1}$. Notice that $V_j := e_0 - e_j$, $j = 1, \dots, n$ is orthogonal to $(1, 1, \dots, 1)$, so $V_1, \dots, V_n$ provide a basis of $\mathcal{L}^\perp$. It is convenient to define

$$\mathcal{T}_{n-1} := \mathcal{V}_n \cap \mathbb{S}^{n-1},$$

where $\mathbb{S}^{n-1}$ is the unit sphere in $\mathcal{L}^\perp = \text{span}\{V_j\}$. The domain $\mathcal{D}_n$ is a double cone over $\mathcal{T}_{n-1}$. More precisely, let N be one of the intersection points in $\mathbb{S}^n \cap \mathcal{L}$ (there are two such points), and let (r, θ) be polar coordinates in $\mathbb{S}^n$, centered at N . Then

$$\mathcal{D}_n = \{(r, \theta) \mid \theta \in \mathcal{T}_{n-1}, 0 \leq r \leq \pi\},$$

where we have identified $\mathbb{S}^{n-1}$ with the unit sphere in the tangent space $T_N \mathbb{S}^n$. In Figure 1 we show $\mathcal{T}_1$ and $\mathcal{D}_2$.

In later sections, we will use a generalization of this type of spherical cone. In general, let Ω be a domain in the equatorial $\mathbb{S}^{n-1}$ of $\mathbb{S}^n$, and let $r_0 \in (0, \pi]$. Then we define the truncated cone

$$\mathcal{TC}(\Omega, r_0) := \{(r, \theta) \mid \theta \in \Omega, 0 \leq r \leq r_0\}.$$

We abbreviate $\mathcal{TC}(\Omega, \pi) = \mathcal{TC}(\Omega)$. In this notation, $\mathcal{D}_n = \mathcal{TC}(\mathcal{T}_{n-1}) = \mathcal{TC}(\mathcal{T}_{n-1}, \pi)$.

The domain $\mathcal{T}_{n-1}$ has symmetry. If we let

$$\mathcal{C}_{n+1}^j = \{(x_0, x_1, \dots, x_n) \mid x_j \geq x_k, j \neq k\},$$

then we see $\mathcal{C}_{n+1} = \mathcal{C}_{n+1}^0$ and $\mathcal{C}_{n+1}^j$ are pairwise congruent. Thus $\mathcal{T}_{n-1}$ is a face of the regular $(n+1)$ -hedral tessellation of the standard $\mathbb{S}^{n-1}$ one obtains by connecting the vertices of a regular $(n+1)$ -simplex with great circle arcs. In particular, one can compute the diameter of $\mathcal{T}_{n-1}$ as the distance from a vertex to the center of the opposite face, which is

$$\delta(n-1) = \arccos\left(-\sqrt{\frac{n-1}{2n}}\right).$$




Figure 1: This figure shows $\mathcal{T}_1$ and $\mathcal{D}_2$.

Moreover, the spherical angle of $\mathcal{T}_{n-1}$ at a vertex is $\mathcal{T}_{n-2}$. So we can construct a succession of comparison domains for $\mathcal{T}_1, \mathcal{T}_2, \dots$ starting with $\mathcal{T}_1$ and using the coning construction described above. To this end, we let

$$\hat{\mathcal{T}}_1 := [0, \frac{2\pi}{3}] = \mathcal{T}_1, \quad \hat{\mathcal{T}}_n := \mathcal{TC}(\hat{\mathcal{T}}_{n-1}, \delta(n)).$$

By induction, $\mathcal{T}_n \subset \hat{\mathcal{T}}_n$, and so $\lambda_1(\mathcal{T}_n) \geq \lambda_1(\hat{\mathcal{T}}_n)$.

2 Separating variables

We discuss two types of separation of variables in this section. The first type is the separation of variables used in [DB] to estimate expected exit times of Brownian motion from Euclidean cones, and the second is the separation of variables one performs to estimate eigenvalues of a spherical domain with a conical structure.

One needs the domains in question to have a certain amount of regularity in order to separate variables. In particular, these domains are piecewise C^1 , satisfy an exterior cone condition, and are well approximated (in terms of Hausdorff distance) by finite volume, C^1 domains. See the introduction to [DB] for a precise statement. It is straight-forward to verify that all of the domains we consider satisfy DeBlassie's hypotheses. We will refer to these domains as *nice*.

2.1 DeBlassie's separations of variables

We first review DeBlassie's [DB] argument. Consider the cone $C \in \mathbb{R}^{n+1}$ over a domain $D \subset \mathbb{S}^n$:

$$C = \{r\theta \mid r > 0, \theta \in D \subset \mathbb{S}^n\}.$$

Let τ_x be the time it takes for a Brownian particle to exit C , with starting position x , and let $u(x, t) = \mathbb{P}(\tau_x > t)$ be the probability that $\tau_x > t$. Then u satisfies the heat equation

$$\begin{aligned} u_t &= \frac{1}{2} \Delta u, & (x, t) &\in C \times [0, \infty); \\ u(x, 0) &= 1, & x &\in \bar{C}; \\ u(x, t) &= 0, & (x, t) &\in \partial C \times (0, \infty). \end{aligned}$$

In polar coordinates (r, θ, t) , the PDE becomes

$$2u_t = u_{rr} + \frac{n}{r}u_r + \frac{1}{r^2}\Delta_{\mathbb{S}^n}u.$$

Moreover, the solution scales as $u(r, \theta, t) = u(\beta r, \theta, \beta^2 t)$, so we can separate variables and look for a solution of the form $u = R(\xi)U(\theta)$, where $\xi = r^2/2t$. Then we get a positive separation constant $\lambda_j(D)$, and

$$\lambda_j(D) = \frac{\Delta_{\mathbb{S}^n} U}{U} = \frac{4\xi^2 \ddot{R} + (4\xi^2 + 2n\xi) \dot{R}}{R}.$$

Letting $R = \xi^a \rho(-\xi)$, this ODE becomes

$$\xi \ddot{\rho} + \left(a + \frac{n+1}{2} - \xi\right) \dot{\rho} - a\rho = 0,$$

which has solutions of the form $\rho(\xi) = f(a/2, a + (n+1)/2, \xi)$, where f is the confluent hypergeometric function (see *e.g.*, [L]). Using f and restricting to $K \subset D$ compact and $T > 0$, one obtains (see [DB]) an expansion for u of the form

$$u(x, t) = \sum_{j=1}^{\infty} B_j f\left(a_j(n), 2a_j(n) + \frac{n+1}{2}, -\frac{|x|^2}{2}\right) U_j\left(\frac{x}{|x|}\right) \left(\frac{|x|^2}{2t}\right)^{-a_j(n)},$$

where U_j is the j th Dirichlet eigenfunction of $\Delta_{\mathbb{S}^n}$ on D and

$$2a_j(n) = \left[\left(\frac{n-1}{2}\right)^2 + \lambda_j(D)\right]^{1/2} - \frac{n-1}{2}.$$

This sum converges uniformly in $K \times [T, \infty)$. The leading term in the expansion is $u(x, t) \sim B_1 U_1(x/|x|)(|x|/(2t))^{-a_1(n)}$, which yields the conclusion of inequality (2).

2.2 Separating variables on the sphere

Next we separate variables to relate the eigenvalues $\lambda_1(\mathcal{D}_n)$ and $\lambda_1(\mathcal{T}_{n-1})$. First recall that we can write the Laplacian for $\mathbb{S}^n$ in polar coordinates as

$$\Delta_{\mathbb{S}^n} u = u_{rr} + (n-1) \cot r u_r + \csc^2 r \Delta_{\theta} u, \quad (3)$$

where Δ_{θ} is the Laplacian on the equatorial $\mathbb{S}^{n-1}$. This lemma gives us a dimension reduction.

Lemma 2 *Let Ω be a nice domain in an equatorial $\mathbb{S}^{n-1}$ with first eigenvalue $\lambda = \lambda_1(\Omega)$, and let $\mathcal{D} = \mathcal{TC}(\Omega)$ be the double cone over Ω . Then the first Dirichlet eigenvalues of $\mathcal{D}$ and Ω are related by*

$$\lambda_1(\mathcal{D}) = \lambda_1(\Omega) - \frac{n-2}{2} + \sqrt{\frac{(n-2)^2}{4} + \lambda_1(\Omega)}. \quad (4)$$

In particular, $\lambda_1(\mathcal{D}) > 2n+2$ whenever $\lambda_1(\Omega) > 2n$.

Proof: Set $u(r, \theta) = R(r)T(\theta)$, where $R(0) = 0 = R(\pi)$ and $T(\theta) = 0$ for $\theta \in \partial\Omega$. Then u is an eigenfunction on $\mathcal{D}$ with eigenvalue μ precisely when

$$T \ddot{R} + (n-1) \cot r T \dot{R} + \csc^2 r R \Delta_{\theta} T = -\mu T R.$$

Separating variables with a positive separation constant λ yields

$$\frac{\sin^2 r \ddot{R} + (n-1) \sin r \cos r \dot{R} + \mu \sin^2 r R}{R} = \lambda = -\frac{\Delta_{\theta} T}{T}.$$

Choosing T to be the first eigenfunction of Ω , we obtain the ODE

$$\sin^2 r \ddot{R} + (n-1) \sin r \cos r \dot{R} + (\mu \sin^2 r - \lambda) R = 0, \quad (5)$$

which has regular singular points at $r = 0, \pi$. If we try a solution of the form $R = \sin^m r$, for some power m , we find

$$\begin{aligned} 0 &= m(m-1) \sin^m r \cos^2 r - m \sin^{m+2} r + m(n-1) \sin^m r \cos^2 r + (\mu \sin^2 r - \lambda) \sin^m r \\ &= \sin^m r [(m^2 + m(n-2) - \lambda) \cos^2 r + (\mu - m - \lambda) \sin^2 r]. \end{aligned}$$

Both coefficients must vanish, so we have $\mu = m + \lambda$ and $\lambda = m^2 + m(n-2)$. Solving for m , we find

$$m = \frac{2-n}{2} + \sqrt{\frac{(2-n)^2}{4} + \lambda}. \quad (6)$$

Next, observe that if $\lambda_1(\Omega) = 2n$ then $\lambda_1(\mathcal{D}) = 2n+2$. Finally, the formula for $\lambda_1(\mathcal{D})$ is monotone increasing in $\lambda_1(\Omega)$, and so $\lambda_1(\mathcal{D}) > 2n+2$ whenever $\lambda_1(\Omega) > 2n$. $\square$

Remark 1 A second solution to equation (5) has the form $\sin^m r \cos r$, where m is again given by equation (6) but

$$\mu' = \lambda + 3m + n.$$

This eigenfunction vanishes on $\{\pi/2\} \times \Omega$, so it corresponds to a higher eigenvalue.

At this point, we can prove that the expected capture time for $n = 1, 2, 3$ predators is infinite. To prove the expected capture time is infinite, by inequality (2) we need to show $\lambda_1(\mathcal{D}_n) \leq 2n+2$, or, equivalently, that $\lambda_1(\mathcal{T}_{n-1}) \leq 2n$. In the case of $n = 1$, we have $\mathcal{D}_1 = [-3\pi/4, \pi/4]$, and so $\lambda_1(\mathcal{D}_1) = 1 < 4$. In the case $n = 2$, we have $\mathcal{T}_1 = [0, 2\pi/3]$, and so $\lambda_1(\mathcal{T}_1) = 9/4 < 4$. We cannot compute $\lambda_1(\mathcal{T}_2)$ so easily, but we can find a test function to show that $\lambda_1(\mathcal{T}_2) < 6$. We shall find $\lambda_1(\mathcal{T}_2)$ numerically in Section 4. Recall the Rayleigh characterization of the first eigenvalue of a domain Ω :

$$\lambda_1(\Omega) = \inf_{f \in H_0^1(\Omega), f \neq 0} \left(\frac{\int_{\Omega} |df|^2}{\int_{\Omega} f^2} \right).$$

To show that $\lambda_1(\mathcal{T}_2) < 6$, it suffices to find $f_0 \in H_0^1(\mathcal{T}_2)$ so that $\int_{\mathcal{T}_2} |df_0|^2 / \int_{\mathcal{T}_2} f_0^2 < 6$. Let

$$f_0(x) = \sin(\text{dist}(x, \partial\mathcal{T}_2)),$$

and observe that $|df_0|^2 = 1 - f_0^2$ off the set of focal points of $\partial\mathcal{T}_2$, which is a set of measure zero. A computation shows

$$\lambda_1(\mathcal{T}_2) \leq \frac{\int_{\mathcal{T}_2} |df_0|^2}{\int_{\mathcal{T}_2} f_0^2} = \frac{2\pi + \sqrt{3}}{\pi - \sqrt{3}} < 6.$$

One can generalize the eigenvalue relationship (4) to spherical cones of the form $\mathcal{TC}(\Omega, r_0)$, for $0 < r_0 < \pi$, using the hypergeometric function

$$f(\alpha, \beta, \gamma, z) = 1 + \frac{\alpha}{\beta} \frac{z}{\gamma} + \frac{\alpha(\alpha+1)}{\beta(\beta+1)} \frac{z^2}{\gamma(\gamma+1)} + \frac{\alpha(\alpha+1)(\alpha+2)}{\beta(\beta+1)(\beta+2)} \frac{z^3}{\gamma(\gamma+1)(\gamma+2)} + \dots \quad (7)$$

Later we will use the following lemma to relate the eigenvalues of $\hat{\mathcal{T}}_{n-1}$ and $\hat{\mathcal{T}}_n$, which we will then estimate to bound the asymptotic decay rates of probabilistic exit times.

Lemma 3 Let Ω be a nice domain in a equatorial $\mathbb{S}^{n-1} \subset \mathbb{S}^n$ with eigenvalue $\lambda = \lambda_1(\Omega)$. Then the first Dirichlet eigenvalues of Ω and $\mathcal{TC}(\Omega, r_0)$ are related by

$$\lambda_1(\mathcal{TC}(\Omega, r_0)) = \mu = \mu(n, \lambda, r_0), \quad (8)$$

where μ is the first Dirichlet eigenvalue of the ODE (5) on $[0, r_0]$. If $r_0 \geq \pi/2$ then μ is the unique zero of $f(\alpha_1, \beta_1, \gamma_1, (1/2)(1 - \cos r_0))$ in $(m + \lambda, 3m + \lambda + n)$, where m is defined in equation (6)

and

$$\begin{aligned}\alpha_1, \beta_1 &= \frac{1 + \sqrt{(n-2)^2 + 4\lambda} \pm \sqrt{(n-1)^2 + 4\mu}}{2} \\ \gamma_1 &= \frac{2 + \sqrt{(n-2)^2 + 4\lambda}}{2}.\end{aligned}$$

If $r_0 \leq \pi/2$ then μ is the unique zero of $f(\alpha_2, \beta_2, \gamma_2, (1/2)(1 - \cos r_0))$ in $(0, n)$, where

$$\alpha_2, \beta_2 = \frac{n-1 \pm \sqrt{(n-1)^2 + 4\mu}}{2}, \quad \gamma_2 = \frac{n}{2},$$

with f defined by equation (7).

Proof: We separate variables and look for a solution of the form $R(r) = \sin^m r u(r)$, with $u(r) \neq 0$ on $[0, r_0)$, but $u(r_0) = 0$. Then equation (5) becomes

$$0 = \sin^{m+2} r \ddot{u} + (2m+n-1) \sin^{m+1} r \cos r \dot{u} + [m(m+n-2) \cos^2 r + (\mu-m) \sin^2 r - \lambda] \sin^m r u.$$

Now let $u(r) = y(x)$, where $x = (1/2)(1 - \cos r)$, which transforms the ODE above into

$$x(1-x)y'' + (m + \frac{1}{2}n - (2m+n)x)y' - (\lambda + m - \mu)y = 0.$$

The solution to this ODE is the hypergeometric function $y(x) = f(\alpha, \beta, \gamma, x)$, with

$$\begin{aligned}\alpha, \beta &= \frac{2m+n-1 \pm \sqrt{(2m+n-1)^2 - 4\lambda - 4m + 4\mu}}{2} \\ \gamma &= \frac{2m+n}{2}.\end{aligned}$$

The lemma follows from taking $R(r; n, \lambda, r_0) = \sin^m r f(\alpha, \beta, \gamma, (1/2)(1 - \cos r))$, where we choose μ so that $R(r_0; n, \lambda, r_0) = 0$. $\square$

One can use this lemma to compute $\lambda_1(\hat{\mathcal{T}}_{n-1})$ iteratively. In this case we start with $\lambda_1(\hat{\mathcal{T}}_1) = 9/4$ and apply equation (8). For the reader's reference, we include a table of $\lambda_1(\hat{\mathcal{T}}_{n-1})$ and the corresponding lower bound for $a(n)$ (see equation (1)). In this table we use equation (4) and the fact that $\mathcal{D}_n \subset \mathcal{TC}(\hat{\mathcal{T}}_{n-1})$ so $\lambda_1(\mathcal{D}_n) \geq \lambda_1(\mathcal{TC}(\hat{\mathcal{T}}_{n-1}))$.

n	$\lambda_1(\hat{\mathcal{T}}_{n-1})$	lower bound for $a(n)$
2	2.25	.75000000
3	5.00463581	.89614957
4	7.884040724	.99030540
5	10.77018488	1.05417466
6	13.6203196	1.09882819

From this table, one can see that $\lambda_1(\hat{\mathcal{T}}_3)$ is slightly less than 8, so $\hat{\mathcal{T}}_3$ comes close to, but is not quite, an effective comparison domain for proving Theorem 1.

3 The eigenvalue estimate

In this section, we complete the proof of Theorem 1. First observe that it suffices to show $\lambda_1(\mathcal{T}_3) > 8$. In fact, it suffices to find an effective lower bound for $\lambda_1(\mathcal{T}_2)$ using the following scheme. We define λ_{cr} by

$$8 = \mu(3, \lambda_{cr}, \delta(3)), \quad \lambda_{cr} \simeq 5.101267527.$$

If we construct a domain $\mathcal{G}_2 \subset \mathbb{S}^2$ such that $\mathcal{T}_2 \subset \mathcal{G}_2$ and $\lambda_1(\mathcal{G}_2) > \lambda_{cr}$, then by domain monotonicity and equation (8),

$$\lambda_1(\mathcal{T}_3) \geq \lambda_1(\mathcal{TC}(\mathcal{T}_2, \delta(3))) \geq \lambda_1(\mathcal{TC}(\mathcal{G}_2, \delta(3))) > 8,$$

completing the proof of Theorem 1. As the last step in our proof, we construct a domain $\mathcal{G}_2 \subset \mathbb{S}^2$ as a perturbation of $\mathcal{T}_2$, such that $\mathcal{T}_2 \subset \mathcal{G}_2$ and $\lambda_1(\mathcal{G}_2) = 5.102 > \lambda_{cr}$.

Figure 2 shows sketches of the domains $\mathcal{T}_2$, $\mathcal{G}_2$ and $\hat{\mathcal{T}}_2$, after stereographic projection to the plane.




Figure 2: The inner dashed curve is $\mathcal{T}_2$, solid curve is $\mathcal{G}_2$, and the outer dashed curve is $\hat{\mathcal{T}}_2$. We generated this figure using the computer program MAPLE.

Previous work of Rayleigh [R] and Pólya–Szegő [PS] motivates us to consider this type of domain perturbation. They studied the eigenvalue of a planar domain which has the form $\{(r, \theta) \mid 0 \leq r \leq c + \epsilon f(\theta)\}$ in polar coordinates, for some small $\epsilon > 0$, and gave an expression for the eigenvalue λ_1 in terms of ϵf . In our case, we fix $\lambda = \lambda_1$ and find a domain $\mathcal{G}_2$ with λ as its first eigenvalue.

Suppose the functions $R(r)$ and $\Theta(\theta)$ satisfy

$$\begin{aligned} \Theta'' + \lambda\Theta &= 0, & 0 \leq \theta \leq \frac{2\pi}{3}; \\ \Theta(0) &= 0, \\ \Theta(2\pi/3) &= 0, \\ \sin^2 r \ddot{R} + \sin r \cos r \dot{R} + (\mu \sin^2 r - \lambda)R &= 0, & 0 \leq r < \pi; \\ R(0) &= 0. \end{aligned}$$

Then $u(r, \theta) = R(r)\Theta(\theta)$ is the first eigenfunction of $\Delta_{\mathbb{S}^2}$ on its nodal domain $\mathcal{G}_2$. By construction, $\Delta_{\mathbb{S}^2}u + \mu u = 0$, so u is an eigenfunction. Also, u does not change sign on its nodal domain, so it must be the first eigenfunction. In polar coordinates, $u(r, 0) = 0 = u(r, 2\pi/3)$. Let $m = \sqrt{\lambda}$ and set $R(r) = \sin^m r u(r)$, so that equation (5) becomes

$$\sin^2(r) \ddot{u} + (1 + 2m) \sin(r) \cos(r) \dot{u} + (\mu - m - \lambda)u = 0.$$

Next we take $\lambda = 9l^2/4$, corresponding to the l^{th} mode of the interval $[0, 2\pi/3]$, and write the solution in terms of the hypergeometric function:

$$u_l(r) = f(3l/2 + .5 \pm \sqrt{1/4 + \mu}, 1 + 3l/2, (1 - \cos r)/2).$$

Finally, we take the $\mu = 5.102$ superposition of the $l = 1, 3$ modes to define

$$\sin^{3/2}(r) \sin(3\theta/2) H(r, \theta) := (\sin r)^{3/2} u_1(r) \sin(3\theta/2) - .0003 (\sin r)^{9/2} u_3(r) \sin(9\theta/4)$$

and let $\mathcal{G}_2$ be the nodal domain of $\sin^{3/2}(r) \sin(3\theta/2) H$. By construction, $\lambda_1(\mathcal{G}_2) = 5.102 > \lambda_{cr}$.

The domains $\mathcal{T}_2$, $\mathcal{G}_2$ and $\hat{\mathcal{T}}_2$ all have three boundary curves, and two of these boundary curves for each domain lie along the great circle arcs $\theta = 0, 2\pi/3$. For convenience, we convert to a planar domain using stereographic projection, with the south pole corresponding to $r = 0$ in polar coordinates on $\mathbb{S}^2$. The relationship between polar coordinates (r, θ) on $\mathbb{S}^2$ and polar coordinates (ρ, θ) on the plane is $\rho = \tan(r/2)$. Under stereographic projection, the great circle arcs $\theta = 0, 2\pi/3$ correspond to rays emanating from the origin at angles $0, 2\pi/3$. The third boundary curve of $\mathcal{T}_2$ is given by the circular arc

$$(\beta(\theta) \cos \theta - \frac{1}{\sqrt{8}})^2 + (\beta(\theta) - \sqrt{\frac{3}{2}})^2 = \frac{3}{2},$$

which we can rewrite as

$$\rho = \beta(\theta) = \frac{\sqrt{2} \cos(\theta - \pi/3) + \sqrt{2 \cos^2(\theta - \pi/3) + 4}}{2}.$$

Thus showing $\mathcal{T}_2 \subset \mathcal{G}_2$ is equivalent to showing $H(r, \theta) > 0$ along the arc given by $(\rho = \beta(\theta), \theta)$ for $0 \leq \theta \leq 2\pi/3$.

One can see from Figure 2 that $H > 0$ along this third boundary component of $\mathcal{T}_2$. One strategy for a rigorous proof that $H > 0$ is the following. We first evaluate H at a point θ_0 on the curve, checking $H > 0$ at $(\rho = \beta(\theta_0), \theta_0)$, and bound the derivative H_θ on an interval containing θ_0 . Our bound $H_\theta \geq M$ gives us a lower bound $H > H(\theta_0) - M(\theta - \theta_0)$. Thus $H > 0$ on a possibly smaller neighborhood of θ_0 . We then repeat this procedure with each endpoint of this (smaller) interval. One can simplify the computations by observing $\mathcal{T}_2$ and $\mathcal{G}_2$ are symmetric about the ray $\theta = \pi/3$. Then it suffices to evaluate H and its derivative at $\theta = 0, 1/2, 2/3, 2\pi/9, \pi/3$. $\square$.

Given our lower bound $\lambda_1(\mathcal{T}_2) \geq \lambda_1(\mathcal{G}_2) = 5.102$, one can use equation (8) to show $\lambda_1(\mathcal{T}_3) \geq 8.00087815$. This in turn gives $\lambda_1(\mathcal{D}_4) \geq 10.001024501$ and $a(4) \geq 1.00007318$. In the same way, the lower bound $\lambda_1(\mathcal{T}_2) \geq 5.102$ gives $a(3) \geq .90671950$.

4 A numerical computation of the first eigenvalue

In this section we describe a numerical computation approximating the eigenvalue $\lambda_1(\mathcal{T}_2)$, which relies on Stenger's sinc-Galerkin scheme [St]. We will show $\lambda_1(\mathcal{T}_2) \approx 5.159 \dots$, and so $a(3) \approx .9128 \dots$. Also, using equation (8), we have $\lambda_1(\mathcal{T}_3) \gtrsim 8.0691$ and so $a(4) \gtrsim 1.0057$.

Given $h > 0$ and a positive integer k , one defines the k th cardinal sinc function of stepsize h as $S(h, k)(hk) = 1$ and if $z \neq hk$,

$$S(h, k)(z) := \frac{\sin(\pi(z - hk)/h)}{\pi(z - hk)/h}. \quad (9)$$

Following Stenger[St], for $z = x + iy$ we shall let the basis functions be $\phi_{jk}(z) = \alpha_j(x) \times \beta_k(y)$, where

$$\alpha_j(x) = S(j, h) \circ \ln\left(\frac{x}{\frac{\pi}{2} - x}\right), \quad \alpha_{n+1}(x) = \sin^2(x) - \sum_{l=-n}^n \sin^2(x_l) \alpha_l(x) \quad (10)$$

for $j = -n, \dots, n$ with sinc points $x_l = \frac{\pi e^{hl}}{2(1 + e^{hl})}$ and

$$\beta_k(x) = S(k, h) \circ \log \sinh y, \quad \beta_{n+1}(y) = \operatorname{sech}(y) - \sum_{l=-n}^n \operatorname{sech}(y_l) \beta_l(y), \quad (11)$$

for $k = -n, \dots, n$ with sinc points $y_l = \sinh^{-1}(e^{hl})$. We let $x_{n+1} = \pi/2$ and $y_{n+1} = 0$. The dimension of the space of functions ϕ_{jk} is $m = (2n+2)^2$. Also $\alpha_{n+1}(\pi/2) = 1$ and $\beta_{n+1}(0) = 1$, but both are zero at the other sinc points because their interior sinc approximations are subtracted. The functions $\phi_{jk} = \alpha_j \times \beta_k$ form a basis of sinc functions when we are working on a rectangular strip.

The most important property of sinc functions is that they are well-suited to approximating integrals in a strip. In particular, one can increase accuracy in a numerical computation by increasing the number of evaluation points, without recomputing grids (as in finite elements). In order to take advantage of this simplicity, we will conformally map a subdomain of $\mathcal{T}_2$ to a half-infinite strip.

Recall that $\mathcal{T}_2$ is an equilateral triangle on $\mathbb{S}^2$ with all its interior angles equal to $2\pi/3$. Let T_1, T_2, T_3 be the vertices of $\mathcal{T}_2$, let S_j be the midpoint of the side opposite T_j , and let F be the center of mass of $\mathcal{T}_2$. Observe that $\mathcal{T}_2$ is invariant under reflection through the lines FT_j , which divide $\mathcal{T}_2$ into six congruent subtriangles (see Figure 3). Thus the first eigenfunction is invariant




Figure 3: The three marked points are F , T_1 and S_3 .

under these reflections, and we can recover it by restricting to the smaller triangle Ω , which has vertices F, T_1, S_3 . The triangle Ω has a right angle at the vertex S_3 , and an angle of $\pi/3$ at the vertices F and T_1 . The first Dirichlet eigenfunction of $\mathcal{T}_2$, restricted to Ω , will have Dirichlet data on the edge T_1S_3 and Neumann data on the edges FT_1 and FS_3 .

We first transform Ω to a planar domain (which we again denote as Ω) using stereographic project, sending S_3 to 0. This transformation sends F to $\frac{1}{2}(\sqrt{6} - \sqrt{2})$ and T_1 to $\frac{i}{2}(\sqrt{6} - \sqrt{2})$. Now Ω is bounded by the two straight line segments joining 0 to $\frac{1}{2}(\sqrt{6} - \sqrt{2})$ and 0 to $\frac{i}{2}(\sqrt{6} - \sqrt{2})$ and the circular arc joining $\frac{1}{2}(\sqrt{6} - \sqrt{2})$ to $\frac{i}{2}(\sqrt{6} - \sqrt{2})$ which makes an angle of $\pi/3$ with the axes. Next we find a conformal transformation f which takes Ω to the half-infinite strip

$$D = \{z \in \mathbb{C} \mid 0 < \Re z < \pi/2, 0 < \Im z\}$$

such that $f(0) = 0$, $f(\frac{i}{2}(\sqrt{6} - \sqrt{2})) = \pi/2$, and $f(\frac{1}{2}(\sqrt{6} - \sqrt{2})) = \infty$. Under this transformation, the first eigenfunction u satisfies

$$\begin{aligned} \Delta^* u + \lambda u &= 0, & \text{if } z \in D, \\ u &= 0, & \text{if } \Re z = 0 \text{ and } \Im z > 0, \\ \frac{\partial u}{\partial n} &= 0, & \text{if } 0 < \Re z < \frac{\pi}{2} \text{ and } \Im z = 0, \text{ or if } \Re z = \frac{\pi}{2} \text{ and } \Im z > 0; \end{aligned}$$

where $\Delta^* = f^* \Delta = \Delta_z$ is the pulled back Laplacian.

There are formulæ for conformally mapping domains bounded by finitely many circular arcs [N] generalizing the Schwarz-Christoffel formula. Following Forsythe [F], the Schwarz triangle mapping $z \in D$ or from $\sin^2 z \in \{\Im z > 0\}$ of the upper halfplane to $w \in \Omega$ is given by

$$\cos^2 z = \frac{(w^4 + 2\sqrt{3}w^2 - 1)^3}{(w^4 - 2\sqrt{3}w^2 - 1)^3}.$$

The group generated by reflections of Ω along its edges in fact tiles the sphere. The inverse function $w \mapsto z^2$ extends to a single sheeted cover and is invariant under the symmetry group, thus is a rational function of the plane. Thus we may compute f . Writing $g(z) = \cos^{2/3} z$,

$$f(z) = \sqrt{\frac{1-g}{\sqrt{3}(1+g) + 2\sqrt{1+g+g^2}}}$$

Pulling back under $w = f(z)$, the conformal weight takes the form

$$\frac{4 \left| \frac{df}{dz} \right|^2}{(1+|f|^2)^2} = \frac{\frac{4}{3} \left| \sqrt{3}(1+g) + 2\sqrt{1+g+g^2} \right|}{|g| \left(\left| \sqrt{3}(1+g) + 2\sqrt{1+g+g^2} \right| + |1-g| \right)^2}.$$

The branch cuts for the square and cube roots may be taken above the negative real axis. Thus $g(D)$ lies in the fourth quadrant so the denominator in f is nonvanishing.

Let $G(z, z')$ denote the Green's function for the problem on D

$$\begin{aligned} \Delta^* u &= g, & \text{if } z \in D, \\ u &= 0, & \text{if } \Re z = 0 \text{ and } \Im z > 0, \\ \frac{\partial u}{\partial n} &= 0, & \text{if } 0 < \Re z < \pi/2 \text{ and } \Im z = 0 \text{ or if } \Re z = \pi/2 \text{ and } \Im z > 0. \end{aligned}$$

The Green's function may be found by the method of images. Thus, denoting $w = \sin z = x + iy$, $w^* = -x + yi$ and $\omega = \sin \zeta$, we get $\bar{w}^* = (\bar{w})^*$. The Green's function is

$$G(z; \zeta) = \frac{1}{2\pi} \left(\ln |w - \omega| - \ln |w^* - \omega| + \ln |\bar{w} - \omega| - \ln |\bar{w}^* - \omega| \right).$$

Pulling back by f , the eigenvalue problem for the triangle may be restated as finding an eigenvalue for the integral operator

$$\frac{1}{\lambda} u(z) = -4 \int_D \frac{G(z, z') |df(z')|^2 u(z') dz'}{(1+|f(z')|^2)^2} =: \mathcal{A}u(z) \quad (12)$$

The key point is to notice that the operator has logarithmic and algebraic singularities at the points $z' = 0, \pi/2$ and $z = z'$. Thus we need to handle these singularities. The solution is zero along the imaginary axis, but free along the other two sides, which may be extended to functions to the plane which have odd reflection symmetry along the imaginary axis and even reflection symmetry along the other sides. We shall approximate $u(z)$ in an m -dimensional space X_m with the same symmetries. Also, noticing that the eigenfunction on $\mathcal{T}_2$ behaves like $1 - \text{dist}^2(z, T_2)$ at the vertex F , we actually have $u \in \text{Lip}(\bar{D})$ and u decays algebraically at ∞ . Choosing a basis $\{\phi_1, \dots, \phi_m\}$ of X_m , we shall compute the matrix of the transformation $A_{\ell k} = \mathcal{P}_\ell \mathcal{A} \phi_k$, whose largest eigenvalue $\mu_m \rightarrow 1/\lambda$ as $m \rightarrow \infty$ and which is an upper bound $\lambda \leq 1/\mu_m$ [St]. The integral operator shall be computed numerically via sinc quadrature, which can handle such mild singularities.

We make the approximation $u(z) \approx \mathcal{P}u(z) = b^{jk} \phi_{jk}(z)$ where the sums are over $j = -n \dots, n+1$ and $k = 0, \dots, n+1$. Here $\phi_{jk} = \alpha_j \times \beta_k$, where α_j and β_k are defined in equations (10) and (11), respectively. We define $b^{jk} = \mathcal{P}_{jk}u = u(x_j + iy_k)$. Thus the approximation $\mathcal{P}u$ is a collocation, as it equals u at the sinc points. Thus, the matrix is approximated by

$$\begin{aligned} A_{jk,pq} &= \int_D G(x_j, y_k, \xi, \eta) \phi_{pq}(\xi, \eta) \Psi(\xi, \eta) d\xi d\eta \\ &\approx \sum_{\iota, \kappa} v_\iota w_\kappa G(x_j, y_k, x_\iota, y_\kappa) \phi_{pq}(x_\iota, y_\kappa) \Psi(x_\iota, y_\kappa) \end{aligned}$$

where $\Psi(\xi, \eta) = 4(1 + |f(\xi + i\eta)|^2)^{-2} |df(\xi + i\eta)|^2$. The approximating sum is carried over $4m$ terms, corresponding to sinc quadratures in the four regions bounded by singularities (*e.g.* in case $0 < x_j < \pi/2$ and $0 < y_k$): $D_I = \{\xi + i\eta : 0 < \xi < x_j, 0 < \eta < y_k\}$, $D_{II} = \{\xi + i\eta : x_j < \xi < \pi/2, 0 < \eta < y_k\}$, $D_{III} = \{\xi + i\eta : 0 < \xi < x_j, y_k < \eta\}$ and $D_{IV} = \{\xi + i\eta : x_j < \xi, y_k < \eta\}$ and where v_ι and w_κ are corresponding weights and x_ι and y_κ are corresponding sinc points.

The computation shows that $\lambda_1(\mathcal{T}_2) \approx 5.159 \dots$ which is above the critical value. This is computational evidence that the exit time has finite expectation. We present a table of a few the computed eigenvalues for approximations in spaces of given dimension, coming from the Sinc-Galerkin collocation scheme described.

Dimension	Eigenvalue Estimate
16	5.948293885960918
100	5.262319373675790
1024	5.153693139833067
2116	5.158585939808193
2304	5.158832705984016
2500	5.158849530710926
2704	5.158968860560663

References

- [BG] M. Bramson and D. Griffeath. *Capture problems for coupled random walks*. in Random Walks, Brownian Motion and Interacting Particle Systems, ed. R. Durrett and H. Kesten, Birkhäuser , 1991.
- [DB] R. D. DeBlassie. *Exit times from cones in $\mathbb{R}^n$ of Brownian motion*. Prob. Theory and Rel. Fields. 74:1–29, 1987.
- [F] A. R. Forsythe. *Theory of Functions of a Complex Variable, 3rd ed.* Dover, 1965, 698–700 (originally published by Cambridge University Press, Cambridge, 1918).
- [L] N. Lebedev, *Special Functions & Their Applications*. Dover, 1972, 260–262. (Originally published by Prentice-Hall, Inc., 1965.)
- [LS] W. Li and Q.-M. Shao. *Capture time of Brownian pursuits*. Prob. Theory and Rel. Fields. 121:30–48, 2001.
- [N] Z. Nehari. *Conformal Mapping*. Dover, 1975 (originally published by Mc Graw-Hill Book Co., Inc., New York, 1952).
- [PS] G. Pólya & G. Szegő. *Isoperimetric Inequalities in Mathematical physics*. Princeton University Press, 1951, 29–30.
- [R] Lord Rayleigh (J. W. Strutt) *The Theory of Sound (2nd. ed.)* Dover, 1945, 336–342 (Originally published by Mcmillan Co., 1894–1896).
- [St] F. Stenger *Numerical Methods Based on Sinc and Analytic Functions* Springer-Verlag, 1993.